

*Disclaimer: This is a machine generated PDF of selected content from our databases. This functionality is provided solely for your convenience and is in no way intended to replace original scanned PDF. Neither Cengage Learning nor its licensors make any representations or warranties with respect to the machine generated PDF. The PDF is automatically generated "AS IS" and "AS AVAILABLE" and are not retained in our systems. CENGAGE LEARNING AND ITS LICENSORS SPECIFICALLY DISCLAIM ANY AND ALL EXPRESS OR IMPLIED WARRANTIES, INCLUDING WITHOUT LIMITATION, ANY WARRANTIES FOR AVAILABILITY, ACCURACY, TIMELINESS, COMPLETENESS, NON-INFRINGEMENT, MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Your use of the machine generated PDF is subject to all use restrictions contained in The Cengage Learning Subscription and License Agreement and/or the Gale OneFile: Health and Medicine Terms and Conditions and by using the machine generated PDF functionality you agree to forgo any and all claims against Cengage Learning or its licensors for your use of the machine generated PDF functionality and any output derived therefrom.*

# Feasibility of Image-Guided Radiotherapy for Elderly Patients with Locally Advanced Rectal Cancer

**Authors:** Nam P. Nguyen, Misty Ceizyk, Jacqueline Vock, Paul Vos, Alexander Chi and Vincent Vinh-Hung

**Date:** Aug. 13, 2013

**From:** PLoS ONE (Vol. 8, Issue 8)

**Publisher:** Public Library of Science

**Document Type:** Article

**Length:** 3,464 words

**DOI:** <http://dx.doi.org/10.1371/journal.pone.0071250>

---

Author(s): Nam P. Nguyen 1,\*, Misty Ceizyk 1, Jacqueline Vock 2, Paul Vos 3, Alexander Chi 4, Vincent Vinh-Hung 5, Judy Pugh 6, Rihan Khan 7, Christina Truong 1, Gabby Albala 1, Angela Locke 1, Ulf Karlsson 8, Steve Gelumauskas 1, Lexie Smith-Raymond 1

## Introduction

Standard of care for locally advanced rectal cancer has been surgery combined with chemoradiation either preoperatively or postoperatively. Preoperative chemoradiotherapy is often preferred because of improved loco-regional control and improved sphincter preservation [1]. The chemotherapy regimen is often based on 5-fluorouracil (5-FU) or capecitabine with similar efficacy when combined with radiotherapy [2]. Grade 3-4 gastrointestinal, urologic, and hematologic toxicities are frequently the limiting factors when chemotherapy is combined with three-dimensional conformal radiotherapy (3D-CRT) because of excessive irradiation of normal pelvic organs [1]-[3]. Given the fear of severe toxicity, elderly patients are often excluded from randomized trials, thus potentially depriving them from a curative treatment. Less aggressive treatment in elderly rectal cancer patients has been reported to increase their cancer-specific mortality even though they may be diagnosed in less advanced stages [4]. New modalities of radiation treatment such as intensity-modulated radiotherapy (IMRT) may potentially decrease the rates of grade 3-4 acute toxicity and improve patient tolerance to chemoradiation by generating a steep dose gradient [5]. Preliminary results with IMRT are encouraging with fewer treatment breaks, less serious toxicities, and less hospitalization in patients with rectal cancer undergoing chemoradiation with various chemotherapy regimens [6]. In addition to reduced toxicity, an excellent pathological response rate and complete resection rate are reported following chemoradiation with IMRT in patients with locally advanced rectal cancer [7]. Thus, the potential advantage in normal tissue sparing that is associated with IMRT may allow elderly patients to receive curative treatment despite the presence of age-associated co-morbidities. Image-guided radiotherapy (IGRT) is a special technique of IMRT delivery combining the steep dose-gradient with accurate daily imaging allowing for precise target radiation dose delivery and further sparing of the small bowels, bladder, and bone marrow [8], [9]. Previously, we reported the feasibility of IGRT to reduce treatment toxicity and improved pathological response in patients with locally advanced rectal cancer undergoing chemoradiation [10]. In the current study, we further assess the efficacy and toxicities of definitive radiotherapy delivered with IGRT in elderly patients receiving curative treatment for locally advanced rectal cancer.

## Materials and Methods

The medical records of 30 patients undergoing neo-adjuvant radiotherapy for locally advanced rectal cancer at the University of Arizona Radiation Oncology department were retrospectively reviewed. The University of Arizona institutional review board

(IRB) approved this retrospective study and waived the requirement for patient consent because of the nature of the study. Locally advanced tumors were defined as T3, T4 tumors based on preoperative ultrasound (US) staging. All patients had a Karnofsky performance status of 70% or higher. All patients had a complete history and physical examination, a digital rectal exam, a CT scan of the chest, abdomen, and pelvis, and an endoluminal US exam. Laboratory tests included a transaminases, alkaline phosphatase, total bilirubin, and carcinoembryonic antigen. Prior to treatment, each patient was simulated in the supine position with a body vacuum bag for treatment immobilization. A computed tomography (CT) scan with and without intravenous (IV) contrast for treatment planning was performed in the treatment position. The abdomen and pelvis were scanned with a slice thickness of 3 mm. Rectal and IV contrast were employed to aid in tumor localization and in identifying grossly enlarged regional lymph node for target volume delineation. Radiotherapy planning was performed on the non-contrast enhanced CT scan to avoid possible interference of contrast density with isodose distribution calculations. Diagnostic positron emission tomography (PET)-CT scan for tumor imaging was also incorporated with CT planning when available. Normal organs at risk (OAR) for complications were outlined for treatment planning (small bowels, bladder, and femoral heads). The gross tumor volume (GTV) was outlined integrating information obtained from the CT scan with IV and rectal contrast, the endoscopic exam, and PET scanning when available. The clinical target volume (CTV) included the rectum, mesorectum, presacral space, and internal iliac nodes. The external iliac nodes were included if there was tumor extension to the vagina, uterus, cervix, prostate, or bladder. The inguinal lymph nodes were treated if there was tumor invasion of the anal canal. The planning target volume (PTV) was generated by isotropically expanding the CTV by a 1-cm margin. The peritoneal cavity was contoured to represent the small bowel volume as it seems to be the most accurate predictor of acute lower gastrointestinal toxicity to pelvic irradiation compared to contouring single bowel loops [11]. An integrated boost technique was used for the IGRT technique to treat the PTV to 45 Gy at 1.8 Gy/fraction and the GTV to 50 Gy at 2 Gy/fraction respectively. Target volume coverage was specified to be at least 95% of the prescribed dose. Dose constraints for normal organs at risk (OAR) for complications were: small-bowel volume receiving 45 Gy (V45) less than 10%, bladder: V45 less than 50%; and femoral head volumes receiving 40 Gy (V40) less than 50%. Twenty-eight patients were treated on a helical tomotherapy unit and 2 patients on a Varian EX unit. Daily MV CT (Tomotherapy) or fusion images (Varian EX) were checked for treatment accuracy. All the radiation constraints were met for tumor coverage and OAR.

## **Chemotherapy**

Concurrent chemotherapy was either intravenous 5-fluorouracil (5FU) or oral capecetabine. Capecetabine was given at an oral dose of 825 mg/m<sup>2</sup> twice daily, 7 days weekly, beginning on the first day of radiotherapy and ending on the last day. 5-FU was administered via continuous intravenous infusion with a portable pump, at 225 mg/m<sup>2</sup> daily during the whole course of radiotherapy. The patients were monitored during treatment with weekly CBC, liver enzymes, electrolytes, BUN, and creatinine. Acute and late treatment toxicity were scored according to the Radiation Therapy Oncology Group (RTOG) scale (<http://ctep.cancer.gov>).

## **Surgery**

Patients underwent surgery 6 to 8 weeks after completion of external beam radiation. All patients underwent a rectal endoscopy prior to surgery to assess tumor response following chemoradiotherapy. The type of surgical procedure was determined by the surgeon based on tumor response. A sphincter-sparing procedure was attempted if tumor shrinkage was deemed sufficient to permit complete resection with negative margins.

## **Pathologic Evaluation**

Resection margins were measured based from the inked surface of the surgical specimen. The circumferential resection margin (CRM) was scored as positive if the tumor was located 1 mm or less from the inked non-peritonealized surface of the specimen. A pathological complete response (pCR) was defined as no residual tumor in the surgical specimen. For patients who did not achieve a pCR, tumor size and depth of invasion were assessed on the surgical specimen and staged according to the TNM system.

## **Statistical Analysis**

Statistical analysis was performed with the log-rank test. A p-value of <0.05 is considered statistically significant. Survival data was analyzed using Kaplan-Meier estimation.

## **Results**

### **Patient Characteristics**

We identified 30 patients with locally advanced adenocarcinoma of the rectum treated with preoperative IGRT chemoradiation at the University of Arizona Radiation Oncology department from 2008 to 2012. Thirteen patients were 70 years or older (elderly) and 17 were less than 70 years (younger), Median age at diagnosis was 77 (range:70-85) and 62 years (range:49-68)

for elderly and younger patients respectively. There were four females in both groups, nine males in the elderly group and 13 males in the younger group. All patients had T3 disease on endoscopic ultrasound. Perirectal lymph nodes were present in three patients, two of the elderly group and one of the younger group. Chemotherapy consisted of 5-FU in 19 patients (7 of the elderly and 12 of the younger group), and capecitabine in 11 patients (6 of the elderly and 5 of the younger group). Table 1 summarizes patient characteristics.

### **Acute Toxicities**

During radiotherapy, one patient in the elderly group developed grade 4 diarrhea secondary to pseudomembranous colitis and required a treatment break of 13 days. Another elderly cancer patient mistook capecitabine for another medication and had an overdose resulting in grade 4 diarrhea and anemia requiring repeated blood transfusion. Her radiotherapy was discontinued after 28 Gy. One patient in the younger group had grade 3 diarrhea. There was no hematologic toxicity in the younger group. One patient in the younger group developed grade 3 diarrhea.

Mean weight loss was 3.7 and 3.9 pounds for elderly and younger patients respectively. No patient in the younger group had a radiotherapy treatment break. Only one patient in the elderly group had a treatment break secondary to his pseudomembranous colitis.

Among the five patients who had chemotherapy protocol violations (dose reduction, delay or discontinuation of chemotherapy), two were in the elderly group and three in the younger group.

Among four patients who did not undergo surgery following chemoradiation, three were in the elderly group and one in the younger group. One elderly patient had surgery aborted because of liver metastases at laparotomy, another one died from cardiac arrhythmia before the scheduled surgery, and the third one declined surgery. The patient in the younger group declined surgery when she was told that she would need an abdomino-pelvic resection because of the tumor location.

All 26 operated patients achieved a complete resection with negative margins. Among the nine patients (30%) who achieved a pCR, three were in the elderly group and the other six were in the younger group.

### **Treatment Outcomes**

After a median follow-up of 42 months (range 3-57), the 3-year survival was estimated to be 90.9% and 87.5% for the elderly and younger patient respectively ( $p = 0.7$ ). No patient who underwent surgery developed a local recurrence. The local recurrence rate was estimated to be 7.6% and 6.6% for the elderly and younger patients respectively ( $p = 0.4$ ). Corresponding numbers for distant metastasis were 17.6% and 15.4% ( $p = 1$ ). Among five patients who developed distant metastases (liver:4, lung and liver:1), two were in the elderly group and three were in the younger group. One patient in the younger group developed a second lung primary and was salvaged with surgery. The two patients who developed long-term complications were in the younger group. One had ischemic colitis and the other one had rectal incontinence. Table 2 summarizes treatment outcomes and toxicities in the two groups.

### **Discussion**

To our knowledge, this is the first study looking at the feasibility of IGRT for elderly patients with locally advanced rectal cancers. Despite a small number of patients, our study suggests that elderly rectal cancer patients tolerate chemoradiation quite well and are able to undergo surgery with excellent loco-regional control because of a high complete resection rate. In the past, there were concerns that elderly rectal cancer patients may not tolerate the combined modality very well because of the expected toxicity of treating a large volume of bowels with conventional chemoradiation technique [12], [13]. Indeed, in studies that reported a high rate of gastrointestinal toxicities and poor tolerance to the combined modality in elderly rectal cancer patients, one-third of the patients were treated with postoperative chemoradiation which has been associated with a higher acute and long-term toxicity compared to preoperative chemoradiation [13], [14]. In addition, most of the treatment breaks were secondary to radiation enteritis associated with excessive bowel irradiation with 3D-CRT [13]. As an illustration, death from myocardial infarction and protocol violations requiring chemotherapy dose reduction and/or radiotherapy interruption secondary to severe toxicity have been reported in 32% of patients with locally advanced cancer treated with capecitabine and oxaliplatin concurrently with 3D-CRT [15]. However, using the same chemotherapy regimen, Sola et al reported that pelvic radiation with IMRT for locally advanced rectal cancer was better tolerated with less interruption of the treatment schedule [7]. Thus, bowel sparing and bone marrow sparing through IMRT may improve patient tolerance to chemoradiation, allowing them to undergo surgery for complete resection and possible cure for their rectal cancer [5], [6]. Image-guided radiotherapy may further improve patient tolerance to chemoradiation because of daily CT imaging allowing for more accurate radiation delivery, and rapid dose fall off comparing to conventional IMRT technique [9], [16]. A simultaneous integrated tumor boost (SIB) delivering a higher radiation dose to the gross tumor while sparing the normal organs may also be achieved to improve local control without excessive increase in acute toxicity [17]. The favorable acute toxicity profile of IGRT may be beneficial to improve elderly rectal cancer patients tolerance to radiation and allow them to have curative resection despite the associated co-morbidity. Indeed, in our study, all patients tolerated chemoradiation quite well and elderly patients fared as well as younger patients with no

significant difference in treatment break or weight loss. Acute grade 3-4 toxicity was acceptable in both groups. Our results compared favorably with other studies using IMRT with chemotherapy for locally advanced rectal cancer. The reported acute grade 3-4 radiation enteritis and hematologic toxicity range from 3% to 24% and 0 to 3% respectively. The percentage of patients who have radiotherapy treatment break range from 11.1% to 16.2% [7], [18]-[20]. Table 3 summarizes acute toxicity in patients with locally advanced rectal cancer treated with IMRT and chemotherapy.

**Table 3. Treatment toxicity reported in studies using IMRT and chemotherapy for the treatment of locally advanced rectal cancer.** [see PDF for image]

Despite the small number of patients, our study highlights the possibility that elderly rectal cancer patients may tolerate chemoradiation better with new techniques of radiotherapy that spare the normal pelvic organs from excessive radiation. Most often, elderly rectal cancer patients are deprived from chemotherapy or radiotherapy because of the age bias [21], [22]. The nihilistic attitude toward elderly patients results in sub-optimal treatment and poor survival (22). Elderly rectal cancer patients who were able to undergo surgery had a significantly better survival compared to the ones who did not have surgery [23]. Image-guided radiotherapy may provide elderly cancer patients a better chance for curative resection and improved survival.

The limitations of the present study include its retrospective nature, the small number of patients, the absence of co-morbidity information, and the relatively short median follow-up. Nevertheless, IGRT may be a new treatment modality to reduce acute toxicity during chemoradiation for locally advanced rectal cancer and improve tolerance to treatment in elderly cancer patients. Further studies with a larger patient population should be performed to investigate the potential of IGRT for curative treatment in elderly rectal cancer patients.

### Conclusion

Image-guided radiotherapy in the setting of chemoradiation is well tolerated in elderly patients with rectal cancer. The potential of IGRT to help elderly rectal cancer patients achieve a curative resection should be investigated in future clinical trials.

### Author Contributions

Conceived and designed the experiments: NPN MC JV PV AC VV JP RK CT GA AL UK SG LSR. Performed the experiments: NPN MC LSR SG. Analyzed the data: NPN MC JV PV AC VV JP RK CT GA AL UK SG LSR. Contributed reagents/materials/analysis tools: AL UK. Wrote the paper: NPN MC JV PV AC VV JP RK CT GA AL UK SG LSR. Literature search: UK PV AC VV. Statistical analysis: PV.

### References

1. Sauer R, Liersch T, Merkel S, Fietkau R, Hohenberger W, et al. (2012) Preoperative versus postoperative chemoradiotherapy for locally advanced rectal cancer: results of the German CAO/ARO/AIO-94 randomized phase III trial after a median follow-up of 11 years. *J Clin Oncol* 30: 1926-1933.
2. Hofheinz RD, Weinz F, Post S, Matzdorff A, Laechelt S, et al. (2012) Chemoradiotherapy with capecitabine versus fluorouracil for locally advanced rectal cancer: a randomized, multicenter, non-inferiority, phase 3 trial. *Lancet Oncol* 13: 579-588.
3. Kalofonos HP, Bamias A, Koutras A, Papakostas P, Basdanis G, et al. (2008) A randomized phase III trial of adjuvant radio-chemotherapy comparing Irinotecan, 5FU and Leucovorin to 5FU and Leucovorin in patients with rectal cancer: A Hellenic Cooperative Oncology Group study. *Eur J Cancer* 44: 1693-1700.
4. Chang GJ, Skibber JM, Feig BW, Rodriguez-Bigas M (2007) Are we undertreating rectal cancer in the elderly? An epidemiologic study. *Ann Surg* 246: 215-221.
5. Arbea L, Ramos LI, Martinez-Monge R, Moreno M, Aristu J (2010) Intensity-modulated radiation therapy vs 3D conformal radiotherapy in locally advanced rectal cancer: dosimetric comparison and clinical implications. *Radiat Oncol* 5: 17.
6. Jabbour SK, Patel S, Herman JM, Wild A, Nagda SN, et al. (2012) Intensity-modulated radiation therapy for rectal carcinoma can reduce treatment breaks and emergency department visits. *Int J Surg Oncol* 890167: 1-7.
7. Arbea L, Martinez-Monge R, Diaz-Gonzales JA, Moreno M, Rodriguez J (2012) et al. (2012) Four-week neoadjuvant intensity-modulated radiation therapy with concurrent capecitabine and oxaliplatin in locally advanced rectal cancer patients: a

validation phase II trial. *Int J Radiat Biol Phys* 83: 587-593.

8. Ippolito E, Mertens I, Haustermans K, Gambacorta MA, Pasini D, et al. (2008) IGRT in rectal cancer. *Acta Oncol* 47: 1317-1324.

9. Beavis AW (2004) Is tomotherapy the future of IMRT? *Br. J Radiol* 77: 285-295.

10. Nguyen NP, Ceizyk M, Almeida F, Chi A, Betz M, et al. (2011) Effectiveness of image-guided radiotherapy for locally advanced rectal cancer. *Ann Surg Oncol* 2011 18: 380-385.

11. Chi A, Nguyen NP, Xu J, Ji M, Tang J, et al. (2012). Correlation of three different approaches of small bowel delineation and acute lower gastrointestinal toxicity in adjuvant pelvic intensity-modulated radiotherapy for endometrial cancer. *Technol Cancer Res Treat* 11, 353-359.

12. Chen RC, Mamon HJ, Ancukiewicz M, Killoran JH, Crowley EM, et al. (2012) Dose-volume effects on patient-reported acute gastrointestinal symptoms during chemoradiation therapy for rectal cancer. *Int J Radiat Oncol Biol Phys* 83: 513-517.

13. Margalit DN, Mamon HJ, Ancukiewicz M, Kobayashi W, Ryan DP, et al. (2011). Tolerability of combined modality therapy for rectal cancer in elderly patients aged 75 years and older. *Int J Radiat Oncol Biol Phys* 81, 735-741.

14. Sauer R, Becker H, Hohenberger W, Rodel C, Wittekind C, et al. (2004). Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 351, 1731-1740.

15. Winkler J, Zipp L, Knoblich J, Zimmermann F (2012) Simultaneous neoadjuvant radiochemotherapy with capecitabine and oxaliplatin for locally advanced rectal cancer. Treatment outcome outside clinical trials. *Strahlenther Onkol* 188: 377-382.

16. Engels B, De Ridder M, Tornel K, Sermeus A, De Coninck P, et al. (2009) Preoperative helical tomotherapy and megavoltage computed tomotherapy for rectal cancer: impact on the irradiated volume of small bowels. *Int J Radiat Oncol Biol Phys* 74: 1476-1480.

17. Engels B, Tournel K, Everaert H, Hoorens A, Sermeus A, et al. (2012) Phase II study of preoperative helical tomotherapy with a simultaneous integrated boost for rectal cancer. *Int J Radiat Oncol Biol Phys* 83: 142-148.

18. Samuelian JM, Callister MD, Ashman JB, Young-Fadok TM, Borad MJ, et al. (2012) Reduced acute bowel toxicity in patients treated with intensity-modulated radiotherapy for rectal cancer. *Int J Radiat Oncol Biol Phys* 82: 1981-1987.

19. Li JL, Ji JF, Cai Y, Li XF, Li YH, et al. (2012) Preoperative concomitant boost intensity-modulated radiotherapy with oral capecitabine in locally advanced mid-low rectal cancer: a phase II trial. *Radiother Oncol* 102: 4-9.

20. Ballonoff A, Kavanagh B, McCarter M, Kane M, Pearlman N, et al. (2008) Preoperative capecitabine and accelerated intensity-modulated radiotherapy in locally advanced rectal cancer: a phase II trial. *Am J Clin Oncol* 31: 264-270.

21. Dale CD (2003) Poor prognosis in elderly patients with cancer: the role of bias and undertreatment. *J Support Oncol* 1: 11-17.

22. Aparicio T, Navazesh A, Boutron I, Bouarioua N, Chosidow D, et al. (2009) Half of elderly patients routinely treated for colorectal cancer receive a sub-standard treatment. *Crit Rev Oncol Hematol* 71: 249-257.

23. Kiran RP, Pokala N, Dudrick SJ (2007) Long-term outcome after operative intervention for rectal cancer in patients aged over 80 years: analysis of 9,501 patients. *Dis Colon Rectum* 50: 604-610.

#### **Author Affiliation:**

**1** Department of Radiation Oncology, University of Arizona, Tucson, Arizona, United States of America, **2** Department of Radiation Oncology, Lindenhofspital, Bern, Switzerland, **3** Department of Biostatistics, East Carolina University, Greenville, North Carolina, United States of America, **4** Department of Radiation Oncology, University of West Virginia, Morgantown, West

Virginia, United States of America, **5** Department of Radiation Oncology, University Hospitals of Geneva, Geneva, Switzerland, **6** Department of Pathology, University of Arizona, Tucson, Arizona, United States of America, **7** Department of Radiology, University of Arizona, Tucson, Arizona, United States of America, **8** Department of Radiation Oncology, Marshfield Clinic, Marshfield, Wisconsin, United States of America

**Corresponding Author:** \* E-mail: NamPhong.Nguyen@yahoo.com

**Editor:** Kevin Robert Kozak, University of Wisconsin School of Medicine and Public Health, United States of America

Article History:

Received Date: 3/4/2013

Accepted Date: 6/26/2013

Published Date: 8/13/2013

**Copyright:** © 2013 Nguyen et al

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The authors have no support or funding to report.

**Competing interests:** The authors have declared that no competing interests exist.

DOI: 10.1371/journal.pone.0071250

Nam P. Nguyen; Misty Ceizyk; Jacqueline Vock; Paul Vos; Alexander Chi; Vincent Vinh-Hung; Judy Pugh; Rihan Khan; Christina Truong; Gabby Albala; Angela Locke; Ulf Karlsson; Steve Gelumbauskas; Lexie Smith-Raymond

**Copyright:** COPYRIGHT 2013 Public Library of Science  
<http://www.plosone.org/static/information>

---

**Source Citation**

Nguyen, Nam P., et al. "Feasibility of Image-Guided Radiotherapy for Elderly Patients with Locally Advanced Rectal Cancer." *PLoS ONE*, vol. 8, no. 8, 2013, p. e71250. *Gale OneFile: Health and Medicine*, <https://link.gale.com/apps/doc/A478298187/HRCA?u=ncliveecu&sid=HRCA&xid=f989362f>. Accessed 27 Mar. 2020.

**Gale Document Number:** GALE|A478298187

---