



Spontaneous bile duct perforation in a neonate

Leslie Hopper^a, Scarlett B. Hao^b, David Rodeberg^b, Shannon Longshore^{b,*}

^a Brody School of Medicine, East Carolina University, USA

^b Vidant Medical Center, East Carolina University, Department of Surgery, Division of Pediatric Surgery, USA



ARTICLE INFO

Keywords:

Bile duct perforation
Bilious ascites
Hepaticojunostomy

ABSTRACT

We present a case of spontaneous biliary perforation (SBP) in a previously healthy 3-week-old boy with worsening scleral icterus, abdominal distension, and acholic stools. Abdominal ultrasound demonstrated moderate ascites without extrahepatic ductal dilatation. The HIDA scan demonstrated accumulation of tracer within the right upper quadrant outside of the liver but not in the bowel, however there was a large amount of tracer activity seen diffusely within the abdomen that appeared to lie within the ascitic fluid and complicated the read of the scan. Intraoperative cholangiogram during an exploratory laparotomy confirmed an intact gallbladder with a mostly contained perforation at the junction of the cystic duct and the common bile duct with no contrast in the duodenum. The patient underwent an uneventful cholecystectomy with a Roux-en-Y hepaticojunostomy and was discharged home three days later.

More than fifty cases of spontaneous biliary perforation (SBP) have been reported in infants and neonates [1,2]. Various etiologies have been suggested, including pancreaticobiliary malunion, ischemia, or an inherent weakness at the junction of the cystic and common hepatic ducts [3–5]. The majority of cases present with acute deterioration or distress at birth, but occasionally these children may have a subacute, indolent presentation. We report on a 3-week-old boy with spontaneous perforation of the bile duct managed with single-stage reconstruction.

1. Case report

A 3-week-old boy, born full term with a benign early postnatal course, presented with worsening jaundice and acholic stools. On examination, positive findings included jaundice and scleral icterus. The abdomen was soft and nondistended with a palpable umbilical hernia and a unilateral hydrocele. Serum liver enzymes were elevated (AST 116 U/L and ALT 88 U/L). Total bilirubin was elevated at 17.8 mg/dL, with a conjugated value of 11.9 mg/dL. Abdominal ultrasound demonstrated a normal liver and gallbladder with no dilation of the intra- or extrahepatic ducts, however there was mild to moderate ascites. A HIDA scan, preceded by a phenobarbital load, demonstrated immediate uptake in the liver, followed by accumulating focal activity in the upper abdomen, thought to be too posterior to represent the gallbladder (Fig. 1).

Of note, the imaging also demonstrated hazy, diffuse activity throughout the abdomen, suspected to be within the ascites, which

made gut activity difficult to confirm. Given the current stability of the child, a laparotomy with intraoperative cholangiogram was performed. Upon entry into the peritoneum, 500 mL of bilious peritoneal fluid was evacuated. An intraoperative cholangiogram demonstrated contrast filling the gallbladder and traversing the hepatic ducts into the liver, however there was significant extravasation of contrast into the left side of the abdomen (Fig. 2).

This finding was confirmed with a second contrast injection after ensuring there was no leak around the cholangiogram catheter. Upon exploration, a small perforation was discovered at the confluence of the cystic and common bile ducts. Additionally, a biloma cavity was noted between the stomach and liver (Fig. 3).

A cholecystectomy and Roux-en-Y hepaticojunostomy was performed, with an anastomotic drain. Post-operatively the neonate tolerated oral feedings with resolution of hyperbilirubinemia (total bilirubin of 3.4 mg/dL from 17.8 mg/dL on admission).

2. Discussion

A recent literature review determined the incidence of SBP to be 1.5 in 1,000,000 live births, with the median age of diagnosis at 2.4 months old [1]. There is no clear etiology for SBP however, three potential etiologies have been proposed. First, a congenital biliary wall weakness due to ischemia of the anterior common bile duct related to a posterolateral arterial supply has been postulated [3,4]. Supporting the theory of congenital wall weakness and ischemia is the observation that the

* Corresponding author. 600 Moye Blvd MA 207, Greenville, NC 27834, USA.
E-mail address: longshores@ecu.edu (S. Longshore).




Fig. 1. HIDA series demonstrating accumulation within the gallbladder, without clear confirmation of duodenal activity due to large amount of diffuse accumulating activity within the abdomen that appeared to lie within the ascitic fluid.

most common site of perforation is at the junction of the cystic duct and common bile duct [1]. A second possible etiology is a pancreaticobiliary malunion near the sphincter of Oddi that would allow reflux of pancreatic juices into the bile duct, leading to acute inflammation and biliary obstruction with increased intraluminal pressure causing perforation [5,6]. Lastly, Moore et al. suggested that SBP seemed to exist on a continuum with choledochal cyst as a form of acquired biliary atresia [7]. This opinion is supported by the observation that SBP seems to occur in children two weeks to five months of age, and perforated choledochal cysts usually present six months to seven years [8,9].

While the presentation of SBP is variable, the most common symptoms are abdominal distension, jaundice, acholic stools, and nonbilious emesis [1]. Evans et al. also noted that hernias and hydroceles were seen in 16% of SBP cases, presumably from increased intra-abdominal pressure following perforation [2]. When presented with a neonate or infant that has jaundice and acholic stools, the surgical differential should include biliary atresia, SBP, and spontaneous perforation of a choledochal cyst. An asymptomatic interval between birth and presentation should decrease the suspicion for biliary atresia, however, SBP and perforation of a choledochal cyst usually present after an asymptomatic interval. It is more difficult to differentiate SBP and perforated choledochal cyst. As noted above, the age of SBP presentation is usually younger than perforated choledochal cysts [8,9].

Imaging is frequently helpful to elucidate the etiology. Abdominal ultrasound has been shown to be a useful diagnostic tool identifying

ascites, ductal dilatation, a fluid collection, or a pseudocyst in cases of SBP [2]. Jeanty et al. noted that RUQ bilomas found during a laparotomy had often been mistaken for an enteric duplication cyst, pancreatic pseudocyst, or choledochal cyst. A HIDA scan is highly sensitive and specific for SBP and is virtually diagnostic when it shows extravasation into the peritoneal cavity with no activity in the duodenum [10]. Over the past several years, magnetic resonance cholangiopancreatography (MRCP) has gained more recognition as a diagnostic tool for SBP. A recent study by Goel et al. stated MRCP may even be better than ultrasound at visualizing a loculated fluid collection or pseudocyst due to its superior soft tissue resolution and multiplanar capability [11].

The optimal management strategy of SBP has been debated. However, nearly all studies agree that an intraoperative cholangiogram should be performed to determine the location of the perforation, to detect any ductal abnormalities, and to exclude distal obstruction [2,12]. If a perforated choledochal cyst is discovered, a single-stage biliary cholecystectomy with a Roux-en-Y hepaticojejunostomy is frequently performed to minimize the risk of developing a cholangiocarcinoma later in life [13]. In the management of SBP, three interventions have been advocated: 1) non-operative management with antibiotics and drainage, percutaneous or ERCP; 2) surgical biliary drainage utilizing external drains, a cholecystostomy tube, or a common duct T-tube; 3) biliary reconstruction via primary repair of the perforation or Roux-en-Y hepaticojejunostomy. Overall, previous




Fig. 2. Intraoperative cholangiogram demonstrating filling of gallbladder (G) and hepatic duct (HD) in addition to significant extravasation (E) into the left side of the abdomen.




Fig. 3. Intraoperative photo of perforation at the confluence of the cystic duct (CyD) and common bile duct (CBD).

studies have favored conservative management with bile drainage (temporary cholecystostomy or T-tube) ± sub-hepatic drainage, resulting in spontaneous closure of perforation [1,13]. Biliary reconstruction can be pursued in patients with poor clinical parameters, pancreaticobiliary malformations, or if distal obstruction is seen on intraoperative cholangiogram [14]. Suture repair of the perforation is not recommended due to concern for post-operative complications including stricture formation, portal vein thrombosis, bile leak, and cholangitis [12].

3. Conclusion

Spontaneous biliary perforation is a rare occurrence in neonates. Abdominal ultrasound in conjunction with a HIDA scan have previously

been shown to be superior for diagnostic imaging when SBP is suspected [2,11]. However, recently it has been suggested that MRCP may offer better visualization than ultrasound when suspecting a fluid collection due to biliary perforation [11]. When present in association with elevated conjugated bilirubin, an intraoperative cholangiogram must be performed in order to establish a diagnosis, confirm perforation, to rule out distal obstruction, and to identify possible pancreaticobiliary malformations. Previous studies have favored external drainage as management of SBP as it is less invasive and requires shorter operative times, even though those same studies have shown similar outcomes between both conservative measures and biliary reconstruction [1,2,12,13]. We chose to manage our patient with a cholecystectomy and Roux-en-Y hepaticojejunostomy due to the fact that our patient had a distal obstruction. However, we propose that the Roux-en-Y

hepaticojejunostomy is an acceptable management option due to surgeon familiarity with the procedure, applicability to all ages, and equivalent outcomes.

Patient consent

Consent to publish the case report was not obtained. This report does not contain any personal information that could lead to the identification of the patient.

Funding

No funding or grant support.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authorship

All authors attest that they meet the current ICMJE criteria for Authorship.

Conflict of interest

The following authors have no financial disclosures: LH, SH, DR, SL.

References

- [1] Jeanty C, Derderian SC, Hirose S, Lee H, Padilla BE. Spontaneous biliary perforation in infancy: management strategies and outcomes. *J Pediatr Surg* 2015;50:1137–41.
- [2] Evans K, Marsden N, Desai A. Spontaneous perforation of the bile duct in infancy and childhood: a systematic review. *J Pediatr Gastroenterol Nutr* 2010;50:677–81.
- [3] Northover JMA, Terblanche J. A new look at the arterial supply of the bile duct in man and its surgical implications. *Br J Surg* 1979;66:379–84.
- [4] Lilly JR, Weintraub WH, Altman RP. Spontaneous perforation of the extrahepatic bile ducts and bile peritonitis in infancy. *Surgery* 1974;75:664–73.
- [5] Hasegawa T, Udatsu Y, Kamiyama M, et al. Does pancreatico-biliary maljunction play a role in spontaneous perforation of the bile duct in children? *Pediatr Surg Int* 2000;16:550–3.
- [6] Ando K, Miyano T, Kohno S, et al. Spontaneous perforation of choledochal cyst: a study of 13 cases. *Eur J Pediatr Surg* 1998;8:23–5.
- [7] Moore TC, Cameron RB. Spontaneous perforation of the extrahepatic biliary tract in infancy and childhood: review of 77 operatively managed cases. *Pediatr Surg Int* 1986;1:206–9.
- [8] Davenport M, Betalli P, D'Antiga L, Cheeseman P, Mieli-Vergani G, Howard ER. The spectrum of surgical jaundice in infancy. *J Pediatr Surg* 2003;38(10):1471–9.
- [9] Chiang L, Chui CH, Low Y, Jacobsen AS. Perforation: a rare complication of choledochal cysts in children. *Pediatr Surg Int* 2011;27:823–7.
- [10] Joodi M, Norouzbeigi N, Ghodsi Rad MA, et al. Spontaneous perforation of common bile duct in a pediatric patient: application of hepatobiliary scintigraphy. *Clin Nucl Med* 2012;37:1007–8.
- [11] Goel P, Jain V, Manchanda V, et al. Spontaneous biliary perforations: an uncommon yet important entity in children. *J Clin Diagn Res* 2013;7:1201–6.
- [12] Patel R, Durell J, Dagash H, et al. Neonatal spontaneous bile duct perforation presenting as giant intraabdominal cyst. *J Pediatr Surg* 2013;1:36–8.
- [13] Babu M, Das N, Das K. Spontaneous infantile choledochal cyst perforation. *Indian J Pediatr* 2007;74:299–300.
- [14] Sahnoun L, Belghith M, Jallouli M, et al. Spontaneous perforation of the extrahepatic bile duct in infancy: report of two cases and literature review. *Eur J Pediatr* 2007;166:173–5.