



Range expansion of a fouling species indirectly impacts local species interactions

Cori J. Speights^{1,2} and Michael W. McCoy²

¹Department of Biological Sciences, Mississippi State University, Starkville, MS, United States of America

²Department of Biology, East Carolina University, Greenville, NC, United States of America

ABSTRACT

We investigated how recent changes in the distribution and abundance of a fouling organism affected the strength of interactions between a commercially important foundation species and a common predator. Increases in the abundance of boring sponges that bioerode the calcified shells of oysters and other shelled organisms have been attributed to increased salinization of estuarine ecosystems. We tested the hypothesis that fouling by boring sponges will change the interaction strength between oysters and a common predator (stone crabs). We generated five oyster density treatments crossed with two sponge treatments (sponge and no sponge). We contrasted the interaction strength between stone crabs and fouled and non-fouled oysters by comparing the parameters of fitted functional response curves based on Rogers random predation model. We found that fouled oysters suffered higher predation from stone crabs, and that the increased predation risk stemmed from a reduction in the handling time needed to consume the fouled oysters. These findings highlight the importance of understanding the effects of abiotic changes on both the composition of ecological communities, and on the strengths of direct and indirect interactions among species. Global climate change is altering local ecosystems in complex ways, and the success of restoration, management, and mitigation strategies for important species requires a better appreciation for how these effects cascade through ecosystems.

Submitted 30 June 2017

Accepted 19 September 2017

Published 19 October 2017

Corresponding author

Cori J. Speights, cjs815@msstate.edu,
cori.speights@gmail.com

Academic editor

Tim Collins

Additional Information and
Declarations can be found on
page 7

DOI 10.7717/peerj.3911

© Copyright

2017 Speights and McCoy

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biodiversity, Conservation Biology, Ecology, Marine Biology

Keywords Boring sponge, Climate change, Fouling, Functional response, Indirect interactions, Predation, Oyster

INTRODUCTION

The strength of interactions between predators and prey can be dependent upon ecological context and a plethora of environmental variables (*Grabowski, 2004; Laudien & Wahl, 1999; Menge, 1995; Wahl, Hay & Enderlein, 1997*). For example interactions with the abiotic environment (e.g., temperature, carbon dioxide, sea level rise) can change activity levels or physiological processes (*Gilman et al., 2010*), and the presence of other organisms can directly or indirectly change the strength of species interactions (*Preisser, Bolnick & Benard, 2005; Werner & Peacor, 2003*). For example, *Schmitt, Osenberg & Bercovitch (1983)* showed that drill holes on kelp snails caused by failed octopus predation attempts resulted in increased barnacle fouling of the snail's shells. The increased fouling increased the

chances of the snails being dislodged from kelp, increasing their exposure to benthic predators by increasing the amount of time spent on the bottom rather than on kelp. Fouling organisms on marine mollusks can also increase susceptibility to predators by compromising the integrity of protective shells (*Duckworth & Peterson, 2013*). These indirect effects, where one species alters the strength of interactions between other species, may become more common and important as species invasions or range expansions resulting from environmental change lead to novel direct and indirect species interactions (*Gilman et al., 2010; Kordas, Harley & O'Connor, 2011; Walther, 2010*).

In marine and estuarine ecosystems, increases in temperature, salinity, and dissolved pCO₂ that are predicted to occur over the next several decades may decrease local habitat quality for some species while facilitating invasions and range expansions for others (*Sorte, Williams & Carlton, 2010; Sunday et al., 2016*). Understanding how changes in biotic and abiotic conditions of ecosystems may change species interactions might be particularly important for foundation species and the communities that depend on their biogenic habitat structures (*Hoegh-Guldberg et al., 2007*). For example, oysters are foundation species in estuaries because their biogenically formed calcium carbonate shells provide habitat structure and refuge that support many other species (*Gutiérrez et al., 2003*). Oysters also provide services such as water filtration that reduces eutrophication, and their reefs provide coastal protection (*Meyer, Townsend & Thayer, 1997; Newell, 2004; Van Wesenbeeck et al., 2013*). Therefore, changes in the distribution of predators or fouling species that affect the health or survival of oysters can have important implications for both oysters and oyster reef communities and the services they provide.

In this study we investigated how the interactions between oysters and a common oyster predator are influenced by a bioeroding sponge which may be expanding its distribution as a result of increasing salinity and temperature in coastal estuaries (*Hong & Shen, 2012; Lindquist, 2011*). Specifically, we investigated how the presence of boring sponges, *Cliona* spp., impact trophic interactions between eastern oysters, *Crassostrea virginica*, and an important native predator, the stone crab *Menippe mercenaria*. While studies have shown stone crabs can have less of an effect on oyster reefs than other mesopredators (e.g., mud crabs) they have recently increased establishment in North Carolina oyster reefs (*Lindquist, 2011; Rindone & Eggleston, 2011*), and we still do not know the magnitude of their effects on oysters interacting with other species, such as sponges. Boring sponges bioerode the calcium carbonate substrates on which they settle (*Duckworth & Peterson, 2013; Fang et al., 2013*). Mollusks that are hosts to boring sponges have weakened shells (*Stefaniak, McAtee & Shulman, 2005*), slower growth, reduced condition, and lower survival than mollusks lacking these bioeroding colonists (*Carroll et al., 2015*). Therefore, we quantified the effects of boring sponges on the interaction strength between stone crabs and fouled and non-fouled oysters. We compared the shape of the crab's functional response to test the hypothesis that the weakened shells of fouled oysters caused by boring sponges will increase the strength of the predator-prey interaction. We focused on the functional response because it is the most direct measure of the interaction strength between predators and prey and it provides a mechanistic link to their population dynamics.

Specifically, to determine the effect of sponges on oyster survival we compared the parameters of type II functional responses (i.e., changes in attack rates or handling times). If sponges are distasteful then crabs will be more likely to avoid foraging on fouled oysters and this effect will be manifested in differences in attack rates. In contrast, if sponges cause changes in shell strength that facilitate crab predation, then we might expect to see shorter handling times and thus higher maximum consumption rates by crabs on fouled oysters.

METHODS

Stone crabs were collected from Middle Marsh in Beaufort, North Carolina (NCDMF Permit No. 706671) and allowed to acclimate in 0.6 m² tanks at the Duke Marine lab for at least 48 h. Each tank received a constant flow of unfiltered seawater and a piece of PVC pipe was provided for refuge. Ten crabs were each wet weighed (g) and the length (mm) of their carapace measured with digital calipers (mean \pm st. error: 93.6 \pm 10.9 g and 64.1 \pm 3.1 mm, respectively). The stone crabs were maintained on a 12 h light/dark cycle and starved for 48 h prior to the beginning of the experiment. Oysters were collected around Morehead City, NC and sorted into two groups: fouled or non-fouled by boring sponge. Oysters of similar sizes (mean \pm st. error: 5.07 \pm 0.07 cm) were used to generate five oyster density treatments of 1,2,4,8, and 22 oysters crossed with two sponge treatments (sponge and no sponge). Oysters were added to each stone crab tank at noon on the day of the experiment. The temperature was recorded in an unused tank at the start of each trial (28.4, 27.7, 25.4, and 24.3 °C, for trials 1, 2, 3, and 4 respectively). The number of oysters eaten was recorded via visual surveys after 24 h and all remaining oysters were removed.

Following each trial, each crab was then fed two oysters a day for three days after which any non-consumed oysters were removed and the crabs were again starved for 48 h and re-randomized for use in another replicate. While not ideal, methods for reusing stone crabs through feeding standardization between trials has been previously reported ([Wong, Peterson & Kay, 2010](#)). In lieu of using new stone crabs for each trial as has been done in previous studies, by using the same crabs, each was influenced by the same background environment before each trial. Additionally, a previous study with rock crabs (*Cancer irroratus*) showed that reused crabs had no change in mussel capture behavior over a three month holding period ([Matheson & Gagnon, 2012](#)). To ensure that any uncertainty due to individual differences among crabs were accounted for we randomly assigned each crab to a sponge \times density treatment for each trial. This distributed any individual crab effect randomly across treatments which minimizes biases in model fits. Two crabs that never consumed oysters in the lab were replaced by new wild-caught crabs for trials. This experiment was replicated four times and oyster collections were made each week to ensure survival of oysters throughout the experiment.

Data were analyzed in the R statistical programming environment ([R Core Team, 2016](#)). Specifically, we fit a Type II functional response curve using Rogers' random predation model ([Juliano, 2001](#); [Rogers, 1972](#)) to quantify predation rates for oysters with and without boring sponge. We used Rogers' formulation because it corrects for prey depletion that occurs as a result of predation over the course of the experiment. The random predator

model predicts the number of prey eaten, N , as:

$$N = N_0(1 - e^{-a(T-hN)}) \quad (1)$$

where T is time, N_0 is the initial prey abundance, h is time spent handling prey and a is the instantaneous attack rate. Rogers' equation can be solved iteratively (Juliano, 2001) as expressed in Eq. (1), however we fit our data to a closed-form solution by expressing Eq. (1) in terms of Lambert's W function (Bolker, 2008; McCoy & Bolker, 2008) so that the number of prey eaten, N , equals:

$$N = N_0 - \frac{W(ahN_0e^{-a(T-hN_0)})}{ah}. \quad (2)$$

Models were fit using the method of maximum likelihood in the *bbmle* package (Bolker & R Development Core Team, 2016) with a binomial error distribution. Specifically, we used a flexible parameter approach to fit (1) a model that estimated attack rates and handling times for the two prey types independently (i.e., a 4 parameter model), which tests the hypothesis that sponges affect both the attack rate and handling times of the prey; (2) a single estimate of attack rate, but separate estimates of handling times (for fouled and non-fouled) (3 parameter model), which test the hypothesis that sponges affect the interaction strength by facilitating crab feeding rates; (3) a model that permitted separate estimates of attack rates (for fouled and non-fouled), but only a single estimate of handling time (3 parameter model), which test the hypothesis that sponges change the likelihood of attack by crabs, and (4) a completely random model that fits only a single estimate of attack rate and handling time (2 parameter model), which serves as our null model. We did not directly measure attack rate or handling time, consequently these parameters were completely estimated from the model. Model fits and inferences about the effects of boring sponges on the interaction between crabs and oysters were made based on sample size-corrected Akaike Information Criterion (AICc).

RESULTS

There was similar support for models 1 and 2 (Table 1), which is interesting given that both of these models allow separate estimates of handling times for crabs eating oysters with and without sponges. This may suggest that sponges are having the largest impacts on crab handling times, which is consistent with previous work indicating that boring sponges weakened mollusks shells (Duckworth & Peterson, 2013). Indeed, handling times (and therefore maximum consumption rates) were approximately 280% longer according to model 1 and 180% longer according to model 2 for crabs eating oysters without sponges relative to oysters with sponges (Table 1). However, the most supported model (model 1 in Table 1) also includes separate estimates of attack rates on oysters with and without boring sponges. While there is no evidence that *Cliona* spp. are unpalatable (Guida, 1976), lower attack rates on fouled oysters may suggest stone crabs have a higher propensity to attack and consume unfouled oysters. Regardless, the differences in attack rates were offset by longer handling times resulting in overall higher consumption of oysters in sponge treatments than in no sponge treatments (Fig. 1).

Table 1 Maximum likelihood results. AICc values for each model. Estimates are presented for all parameters (α = attack rate and h = handling time) allowed to vary by treatment in a model (95% confidence intervals are presented underneath each estimate). With few observations (nobs = 38), corrected AIC (AICc) was used instead of AIC.

Model	Parameters	dAICc	df	Weight	No sponge	α	Sponge	No sponge	h	Sponge
1	a^*h	0	4	0.597	4.079 (1.457, 6.701)		1.970 (-3.416, 7.357)	0.112 (0.075, 0.149)		0.039 (-0.043, 0.123)
2	h	1.1	3	0.352		2.534 (1.660, 3.408)		0.093 (0.063, 0.123)		0.051 (-0.011, 0.113)
3	a	5.5	2	0.038	2.414 (1.140, 3.688)		1.97 (0.060, 5.353)		0.072 (0.048, 0.095)	
4	1	7.7	3	0.013		2.62 (1.690, 3.551)			0.074 (0.052, 0.096)	




Figure 1 Prey consumed over 24 h. Amount of prey consumed by predators over a 24 h period using five increasing densities. Lines represent oysters (*Crassostrea virginica*) with sponges (*Cliona* spp.) (black) and oysters without sponges (gray), with standard error bars for each point ($n = 4$ trials). Attack rates and handling times used for each line were obtained from model 1 (see Table 1).

Full-size DOI: 10.7717/peerj.3911/fig-1

DISCUSSION

We investigated how a fouling species that has expanded its range may be indirectly impacting the eastern oyster. Our results show that the presence of fouling from boring sponges will make oysters more susceptible to predation by crabs and likely other shell-crushing predators. One potential mechanism that we present here is a decrease in predator handling time for oysters with sponges compared to those without sponges. Indeed, boring sponges (*Cliona celata*) have been shown to weaken scallop shells by as much as 28% (Duckworth & Peterson, 2013). However, other studies have suggested that infestation by boring sponges did not impact stone crab handling times (Coleman, 2014). However, this difference may be due in part to difference in the sizes of the stone crabs used in the two studies (mean carapace from Coleman 2014 = 98.5 and current study 64.1 mm), such that defenses in shell strength were only evident for smaller crabs.

Understanding the potential effects of changes in species ranges and interactions are especially important for foundation species that provide structure that serves as primary habitat for communities of other species (Dayton, 1973). Specifically, global environmental

change can facilitate species range expansions and alter local trophic interactions (Walther *et al.*, 2002), which is critical information for mitigating and managing affected ecosystems. Boring sponges, *Cliona* spp., are experiencing range expansions potentially as a result of increased salinization of some estuarine ecosystems (Dunn, Eggleston & Lindquist, 2014; Lindquist, 2011). Boring sponges are generally found in areas with >15 ppt salinity (Carver, Thériault & Mallet, 2010; Hopkins, 1962; Lindquist, 2011), and they are being documented in increasing abundances further up-estuary as sea level and salinity rise (Hong & Shen, 2012; Lindquist, 2011).

Therefore, changes in the range of a fouling species in response to changes in habitat characteristics or climate change can have indirect consequences on the trophic interactions between important species (such as oysters and corals) and their natural enemies. Indeed, studies have shown that boring sponges are not impacted by high water temperatures or decreases in pH (due to an increase in atmospheric CO₂) but instead these factors increase shell boring rates (Duckworth & Peterson, 2013). In addition, documented increases in the stone crab's northward expansion along with boring sponges could enhance potential negative impacts to oyster fisheries in states such as Virginia and North Carolina. Overall, understanding how increases in predation risk as a result of fouling by boring sponges works in concert with other effects of global climate change (i.e., sea level rise, ocean acidification and increasing salinity) will have important implications for managing foundation species and the services they provide through fisheries, coastal protection, and ecosystem engineering.

This study highlights the need to consider how indirect biotic interactions can alter the interaction strengths between predators and prey. Indeed, boring sponges alone have modest impacts on oyster fitness. However, changes in the distribution and abundance of sponges, increases in boring efficiency, and the interactions between boring sponges and other species can lead to strong negative impacts on oysters and oyster reef communities. Such context dependent and indirect effects must be considered in future restoration and management aimed at recovering already heavily damaged oyster reef ecosystems (Beck *et al.*, 2011; D'Anna, 2016).

ACKNOWLEDGEMENTS

We would like to thank Brian Silliman, the members of the Silliman Lab at the Duke Marine Lab, and lab assistants Thomas Guryan and Erin Tomaras for providing lab space and experimental assistance. In addition, we would like to thank David Kimmel, April Blakeslee, Krista McCoy, and the McCoy Labs for constructive comments on this research.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research was supported by East Carolina University and in part by a National Science Foundation Award (#1556743) to Michael McCoy. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
National Science Foundation Award: #1556743.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Cori J. Speights conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Michael W. McCoy conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

For the collection of crabs, we had a collection permit approved by the North Carolina Division of Marine Fisheries. Field study approval number: 706671.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3911#supplemental-information>.

REFERENCES

- Beck MW, Brumbaugh RD, Airoidi L, Carranza A, Coen LD, Crawford C, Defeo O, Edgar GJ, Hancock B, Kay MC. 2011. Oyster reefs at risk and recommendations for conservation, restoration, and management. *Bioscience* 61:107–116 DOI 10.1525/bio.2011.61.2.5.
- Bolker BM. 2008. *Ecological models and data in R*. Princeton, NJ: Princeton University Press.
- Bolker B, R Development Core Team. 2016. bbmle: tools for general maximum likelihood estimation. R package version 1.0.18.
- Carroll JM, Shaughnessy KA, Diedrich GA, Finelli CM. 2015. Are oysters being bored to death? Influence of *Cliona celata* on *Crassostrea virginica* condition, growth and survival. *Diseases of Aquatic Organisms* 117:31–44 DOI 10.3354/dao02928.
- Carver CE, Thériault I, Mallet AL. 2010. Infection of cultured eastern oysters *Crassostrea virginica* by the boring sponge *Cliona celata*, with emphasis on sponge life history and mitigation strategies. *Journal of Shellfish Research* 29:905–915 DOI 10.2983/035.029.0423.
- Coleman SE. 2014. The effects of boring sponge on oyster soft tissue, shell integrity, and predator-related mortality. MSc thesis, University of North Carolina, Chapel Hill, NC.

- D'Anna LM. 2016.** Concern is in the eye of the stakeholder: heterogeneous assessments of the threats to oyster survival and restoration in North Carolina. *Society & Natural Resources* **29**:131–147 DOI [10.1080/08941920.2015.1058998](https://doi.org/10.1080/08941920.2015.1058998).
- Dayton PK. 1973.** Toward an understanding of community resilience and the potential effects of enrichments to the benthos at McMurdo Sound, Antarctica. In: Parker BC, ed. *Proceedings of the colloquium on conservation problems in Antarctica*. Lawrence: Allen Press.
- Duckworth A, Peterson B. 2013.** Effects of seawater temperature and pH on the boring rates of the sponge *Cliona celata* in scallop shells. *Marine Biology* **160**:27–35 DOI [10.1007/s00227-012-2053-z](https://doi.org/10.1007/s00227-012-2053-z).
- Dunn RP, Eggleston DB, Lindquist N. 2014.** Oyster-sponge interactions and bioerosion of reef-building substrate materials: implications for oyster restoration. *Journal of Shellfish Research* **33**:727–738 DOI [10.2983/035.033.0307](https://doi.org/10.2983/035.033.0307).
- Fang JK, Mello-Athayde MA, Schönberg CH, Kline DI, Hoegh-Guldberg O, Dove S. 2013.** Sponge biomass and bioerosion rates increase under ocean warming and acidification. *Global Change Biology* **19**:3581–3591 DOI [10.1111/gcb.12334](https://doi.org/10.1111/gcb.12334).
- Gilman SE, Urban MC, Tewksbury J, Gilchrist GW, Holt RD. 2010.** A framework for community interactions under climate change. *Trends in Ecology & Evolution* **25**:325–331 DOI [10.1016/j.tree.2010.03.002](https://doi.org/10.1016/j.tree.2010.03.002).
- Grabowski JH. 2004.** Habitat complexity disrupts predator–prey interactions but not the trophic cascade on oyster reefs. *Ecology* **85**:995–1004 DOI [10.1890/03-0067](https://doi.org/10.1890/03-0067).
- Guida VG. 1976.** Sponge predation in the oyster reef community as demonstrated with *Cliona celata* Grant. *Journal of Experimental Marine Biology and Ecology* **25**:109–122 DOI [10.1016/0022-0981\(76\)90012-5](https://doi.org/10.1016/0022-0981(76)90012-5).
- Gutiérrez JL, Jones CG, Strayer DL, Iribarne OO. 2003.** Mollusks as ecosystem engineers: the role of shell production in aquatic habitats. *Oikos* **101**:79–90 DOI [10.1034/j.1600-0706.2003.12322.x](https://doi.org/10.1034/j.1600-0706.2003.12322.x).
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K. 2007.** Coral reefs under rapid climate change and ocean acidification. *Science* **318**:1737–1742 DOI [10.1126/science.1152509](https://doi.org/10.1126/science.1152509).
- Hong B, Shen J. 2012.** Responses of estuarine salinity and transport processes to potential future sea-level rise in the Chesapeake Bay. *Estuarine, Coastal and Shelf Science* **104**:33–45 DOI [10.1016/j.ecss.2012.03.014](https://doi.org/10.1016/j.ecss.2012.03.014).
- Hopkins SH. 1962.** Distribution of species of *Cliona* (boring sponge) on the Eastern Shore of Virginia in relation to salinity. *Chesapeake Science* **3**:121–124 DOI [10.2307/1351224](https://doi.org/10.2307/1351224).
- Juliano S. 2001.** Nonlinear curve fitting: predation and functional response curves. *Design and Analysis of Ecological Experiments* **2**:178–196.
- Kordas RL, Harley CD, O'Connor MI. 2011.** Community ecology in a warming world: the influence of temperature on interspecific interactions in marine systems. *Journal of Experimental Marine Biology and Ecology* **400**:218–226 DOI [10.1016/j.jembe.2011.02.029](https://doi.org/10.1016/j.jembe.2011.02.029).

- Laudien J, Wahl M. 1999.** Indirect effects of epibiosis on host mortality: seastar predation on differently fouled mussels. *Marine Ecology* **20**:35–47
DOI [10.1046/j.1439-0485.1999.00063.x](https://doi.org/10.1046/j.1439-0485.1999.00063.x).
- Lindquist N. 2011.** Quantifying boring sponge abundance, biomass and bioerosion rates in North Carolina oyster reefs. North Carolina Sea Grant Fishery Resource Grant Final Report. Available at <https://uncch.pure.elsevier.com/en/projects/quantification-of-boring-sponge-abundance-and-biomass-in-north-ca>.
- Matheson K, Gagnon P. 2012.** Temperature mediates non-competitive foraging in indigenous rock (*Cancer irroratus* Say) and recently introduced green (*Carcinus maenas* L.) crabs from Newfoundland and Labrador. *Journal of Experimental Marine Biology and Ecology* **414**:6–18 DOI [10.1016/j.jembe.2012.01.006](https://doi.org/10.1016/j.jembe.2012.01.006).
- McCoy MW, Bolker BM. 2008.** Trait-mediated interactions: influence of prey size, density and experience. *Journal of Animal Ecology* **77**:478–486
DOI [10.1111/j.1365-2656.2008.01372.x](https://doi.org/10.1111/j.1365-2656.2008.01372.x).
- Menge BA. 1995.** Indirect effects in marine rocky intertidal interaction webs: patterns and importance. *Ecological Monographs* **65**:21–74 DOI [10.2307/2937158](https://doi.org/10.2307/2937158).
- Meyer DL, Townsend EC, Thayer GW. 1997.** Stabilization and erosion control value of oyster cultch for intertidal marsh. *Restoration Ecology* **5**:93–99
DOI [10.1046/j.1526-100X.1997.09710.x](https://doi.org/10.1046/j.1526-100X.1997.09710.x).
- Newell RI. 2004.** Ecosystem influences of natural and cultivated populations of suspension-feeding bivalve molluscs: a review. *Journal of Shellfish Research* **23**:51–62.
- Preisser EL, Bolnick DI, Benard MF. 2005.** Scared to death? The effects of intimidation and consumption in predator–prey interactions. *Ecology* **86**:501–509
DOI [10.1890/04-0719](https://doi.org/10.1890/04-0719).
- R Core Team. 2016.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Rindone RR, Eggleston DB. 2011.** Predator–prey dynamics between recently established stone crabs (*Menippe* spp.) and oyster prey (*Crassostrea virginica*). *Journal of Experimental Marine Biology and Ecology* **407**:216–225 DOI [10.1016/j.jembe.2011.06.018](https://doi.org/10.1016/j.jembe.2011.06.018).
- Rogers D. 1972.** Random search and insect population models. *The Journal of Animal Ecology* **41**:369–383.
- Schmitt RJ, Osenberg CW, Bercovitch MG. 1983.** Mechanisms and consequences of shell fouling in the kelp snail, *Norrisia norrisi* (Sowerby) (Trochidae): indirect effects of octopus drilling. *Journal of Experimental Marine Biology and Ecology* **69**:267–281
DOI [10.1016/0022-0981\(83\)90074-6](https://doi.org/10.1016/0022-0981(83)90074-6).
- Sorte CJ, Williams SL, Carlton JT. 2010.** Marine range shifts and species introductions: comparative spread rates and community impacts. *Global Ecology and Biogeography* **19**:303–316 DOI [10.1111/j.1466-8238.2009.00519.x](https://doi.org/10.1111/j.1466-8238.2009.00519.x).
- Stefaniak LM, McAtee J, Shulman MJ. 2005.** The costs of being bored: effects of a clonoid sponge on the gastropod *Littorina littorea* (L). *Journal of Experimental Marine Biology and Ecology* **327**:103–114 DOI [10.1016/j.jembe.2005.06.007](https://doi.org/10.1016/j.jembe.2005.06.007).

- Sunday JM, Fabricius KE, Kroeker KJ, Anderson KM, Brown NE, Barry JP, Connell SD, Dupont S, Gaylord B, Hall-Spencer JM. 2016.** Ocean acidification can mediate biodiversity shifts by changing biogenic habitat. *Nature Climate Change* 7:81–85.
- Van Wesenbeeck BK, Griffin JN, van Koningsveld M, Gedan KB, McCoy MW, Silliman BR. 2013.** Nature-based coastal defenses: can biodiversity help? In: Levin S, ed. *Encyclopedia of biodiversity*. Waltham: Academic Press, 451–458.
- Wahl M, Hay M, Enderlein P. 1997.** Effects of epibiosis on consumer–prey interactions. *Hydrobiologia* 355:49–59 DOI [10.1023/A:1003054802699](https://doi.org/10.1023/A:1003054802699).
- Walther G-R. 2010.** Community and ecosystem responses to recent climate change. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365:2019–2024 DOI [10.1098/rstb.2010.0021](https://doi.org/10.1098/rstb.2010.0021).
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJ, Fromentin J-M, Hoegh-Guldberg O, Bairlein F. 2002.** Ecological responses to recent climate change. *Nature* 416:389–395 DOI [10.1038/416389a](https://doi.org/10.1038/416389a).
- Werner EE, Peacor SD. 2003.** A review of trait-mediated indirect interactions in ecological communities. *Ecology* 84:1083–1100 DOI [10.1890/0012-9658\(2003\)084\[1083:AROTII\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[1083:AROTII]2.0.CO;2).
- Wong MC, Peterson CH, Kay J. 2010.** Prey size selection and bottom type influence multiple predator effects in a crab–bivalve system. *Marine Ecology Progress Series* 409:143–156 DOI [10.3354/meps08621](https://doi.org/10.3354/meps08621).