



Published in final edited form as:

*J Sleep Res.* 2019 August ; 28(4): e12810. doi:10.1111/jsr.12810.

## Cognitive performance in patients with Implantable Cardioverter Defibrillators (ICDs): Associations with objective sleep duration, age, and anxiety

Ashley F. Curtis, PhD<sup>1</sup>, Alicia J. Roth, PhD<sup>2</sup>, Samuel F. Sears, PhD<sup>3</sup>, Jamie B. Conti, MD<sup>4</sup>, Richard B. Berry, MD<sup>5</sup>, Joseph M. Dzierzewski, PhD<sup>6</sup>, Himangshu Rathinakumar, MBBS<sup>1</sup>, and Christina S. McCrae, PhD<sup>1</sup>

<sup>1</sup>Department of Psychiatry, University of Missouri. Columbia, MO, USA

<sup>2</sup>Department of Psychiatry and Psychology, Cleveland Clinic, Cleveland, OH, USA

<sup>3</sup>Departments of Psychology and Cardiovascular Sciences, East Carolina University, Greenville, NC, USA

<sup>4</sup>Division of Cardiovascular Medicine, University of Florida, Gainesville, FL, USA

<sup>5</sup>College of Medicine, University of Florida, Gainesville, FL, USA

<sup>6</sup>Department of Psychology, Virginia Commonwealth University, Richmond, VA, USA

### Summary

Sleep disturbance and anxiety are highly prevalent in patients with implantable cardiac defibrillators (ICDs). There is limited research, however, on the associations between cognitive performance and sleep parameters, age, and anxiety. Forty-one patients with ICDs and self-reported sleep disturbance completed 14 days of actigraphy ( $M_{age}=60.3$ ,  $SD=12.3$ ) measuring total sleep time (TST), and a computerized cognitive test battery measuring processing speed/attention (i.e., simple reaction time, symbol digit modality task or SDMT) and executive function (i.e., flanker task, letter series task, N-back task). Multiple regressions determined whether independent effects of TST, age, and anxiety, as well as interactive effects of TST and age predicted cognitive performance. TST predicted performance on two tasks of executive function (i.e., letter series, N-back task), as well as an attentional vigilance/processing speed task (i.e., SDMT), and this did not depend on patient age. On letter series, N-back, and SDMT, longer TST predicted better performance. Increasing age predicted worse performance on SDMT and flanker tasks. No other predictors were associated with task performance. Results show that sleep duration, not anxiety, may be an important predictor of higher order cognitive functioning and lower order tasks measuring processing speed and attention in ICD patients, with longer sleep duration showing greater benefit to performance.

**Corresponding Author:** Christina S. McCrae, PhD; Professor; Department of Psychiatry, University of Missouri-Columbia, 1 Hospital Drive, Columbia, MO 65212; Ph +1 573-882-0982; Fax +1 573-884-1070; mccraec@health.missouri.edu.

**Author Contributions:** A.F.C and A.J.R drafted the manuscript. A.F.C. conducted analyses. J.M.D. collected data. C.S.M. and J.M.D. drafted and carried out protocol. All authors revised draft.

**Conflicts of Interest:** S.F.S has research grants from Medtronic and Zoll Medical, has received honorarium from Medtronic, Boston Scientific, Zoll Medical, and Abbott/St Jude Medical in the last year, and is a consultant for Abbott/St Jude Medical.

## Keywords

cardiovascular disease; aging; sleep disturbance; cognitive performance

---

## Introduction

Cardiovascular disease is the leading cause of death in the United States, accounting for approximately 500,000 deaths annually (Heron, 2017). Implantable cardioverter defibrillators (ICDs) are widely used in patients at high risk for sudden death or in those who have had a previous ventricular arrhythmia (Kusumoto et al., 2014). The ICD provides both low-energy pacing and high energy shock to terminate ventricular arrhythmias. These shocks are known to be uncomfortable or painful to the patient (Dornelas and Sears, in press). Up to 67% of patients with ICDs experience disturbed sleep (Habibovi et al., 2018, Berg et al., 2012), and ICD use has also been associated with increased anxiety (Bilge et al., 2006, Sola and Bostwick, 2005, Pedersen et al., 2011, Sears et al., 2011). Research also shows that patients with ICDs experience self-reported cognitive disturbance (Kamphuis et al., 2004), as well as objective cognitive deficits across a wide range of tasks (Hałas et al., 2014), which is exacerbated with increasing age (Hałas et al., 2014). However, the effects of disturbed sleep and anxiety on cognitive functioning in ICD patients has not been established. Thus, examinations of the impact of sleep and the potential moderating influence of age on cognitive performance of ICD patients is warranted (Moulaert et al., 2010).

Although sleep parameters, particularly sleep duration or total sleep time (TST), have been examined in other adult populations, there is a paucity of research documenting effects of disturbed sleep in ICD patients. For instance, self-reported TST (i.e., <6 hours) has been associated with worse performance on tests measuring higher order or executive function in younger adults with insomnia (Khasawneh et al., 2018), processing speed and executive function in middle-aged adults with insomnia (Fernandez-Mendoza et al., 2010), and worse global cognitive performance in older adults (Lo et al., 2014). Other research has found that objectively measured (i.e., via actigraphy) TST is associated with worse global cognitive performance in older adults (Blackwell et al., 2011). Given findings of associations between sleep and cognitive performance, and the well-known finding that cognitive performance declines with increasing age (Salthouse, 2009), in the present study we specifically examined how TST and age potentially interact in their prediction of cognitive performance in patients with ICDs. Such evidence could inform behavioral treatments aimed at improving primary sleep and secondary outcomes such as cognitive performance in these patients, with the goal of ultimately improving daytime functioning and increasing quality of life. Additionally, given the high prevalence of anxiety in ICD patients (Bilge et al., 2006) and the research documenting that greater anxiety can impair performance on tests measuring executive abilities (Derakshan and Eysenck, 2009), we also assessed the impact of self-reported general anxiety on cognitive function.

The goal of this study was to examine associations between objective sleep duration and anxiety on cognitive functioning in patients with an ICD. We also examined whether sleep duration interacted with age in these associations with cognition. We predicted that longer

TST would be associated with better cognitive performance on tasks requiring higher order cognitive functioning (i.e., executive functioning tasks measuring reasoning, inhibition and working memory), but would not affect performance on tasks requiring lower level cognitive processing (i.e., sustained attention/processing speed). We also predicted that these associations would be stronger with increasing age. Finally, we predicted that greater anxiety would be independently associated with worse cognitive performance on tasks requiring higher order cognition.

## Methods

### Participants

Participants were recruited as part of a larger randomized controlled trial ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02232204) Identifier: NCT02232204) through physician referral from a university-based cardiac clinic. Participants were included if they had: 1) an implanted ICD device and 2) reported difficulty sleeping via a screening interview (i.e., responded “yes” to the question “are you currently experiencing difficulty with your sleep?”). Participants were excluded if they demonstrated severe cognitive impairment [Mini Mental Status Exam scores of <23 (if participants had 9<sup>th</sup> grade education or higher) or <17 (if participants had less than 9<sup>th</sup> grade education)], were shift workers, or did not complete 14 days of actigraphy/cognitive tests. Participants completed a clinical interview, where we obtained: demographics, cardiac-specific health history, and current use of prescription sleep medications. A formal review of medical records was conducted prior to the baseline visit to verify cardiovascular-specific history (i.e., date of ICD implantation, cardiac etiology), as well as information regarding premorbid psychiatric diagnoses (e.g., depression, anxiety). The University of Florida Institutional review board approved all study procedures, and participants provided informed consent.

### Sleep Measures

Sleep was assessed through actigraphy. Participants wore an actigraphy watch (Actiwatch 2, Phillips Respironics, Bend, OR) on their nondominant wrist 24 hours a day for the 14 days of baseline. Actigraphy is highly correlated with polysomnography on measures of TST in individuals with insomnia, with moderate correlations for nighttime wake variables (.30 for sleep onset latency, SOL; .48 for wake after sleep onset, WASO) (Lichstein et al., 2006). The Actiwatch 2 records data on gross motor activity using a solid-state, piezo-electric accelerometer. The accelerometer continually measures the intensity and frequency of wrist movement at a sampling rate of 32 cycles/second. The sum of all wrist movements within 30 seconds is recorded as an activity count. Activity counts were analyzed using Actiware Sleep Analysis Software v.5.3.2, which classifies each 30-second epoch as sleep or wake using validated algorithms (Oakley, 1997). The high sensitivity setting was used because it provides good correlation with PSG for TST (.70) in individuals with insomnia (Lichstein et al., 2006, McCrae et al., 2005). As part of the parent clinical trial procedures, participants also completed daily sleep diaries (Lichstein et al., 1999). Bed and wake times reported on sleep diaries that were recorded concurrent with actigraphy measurement were inserted into corresponding actigraph days to represent the time-in-bed period. Actiware determined the start of sleep by searching this time-in-bed period for the first 10-minute interval during which no more than 1 epoch was scored as awake. Similarly, sleep end was signified by the

last 10-minute interval containing no more than 1 wake-state epoch. The average TST value across 14 days of baseline was the sleep parameter of interest in the present study.

Etiology of self-reported sleep disturbance was determined by a board certified sleep physician (R.B.B) and sleep psychologist (C.S.M.), based on an in-depth clinical interview, 14 days of sleep diaries, and one night of ambulatory polysomnography (PSG) testing. Sleep apnea and periodic leg movement disorder (PLMD) were diagnosed in accordance with criteria of the International Classification of Sleep Disorders-3rd ed. (AASM., 2014), and insomnia criteria were consistent with DSM-5 criteria (Association, 2013) and Research Diagnostic Criteria (RDC) (Edinger et al., 2004). Insomnia was diagnosed if participants reported: 1) >30 minutes of SOL (Lichstein et al., 2003), WASO, or waking up too early; 2) sleep that is chronically non-restorative or poor in quality; 3) items 1 and 2 occur despite adequate opportunity/circumstances for sleep; 4) insomnia present 3+ nights/week for more than 1 month; 5) 1+ sleep-related daytime impairment (e.g., fatigue, cognitive disruption, social/vocational/educational disruption, mood disturbance/irritability, daytime sleepiness, motivation/energy/initiative reduction, proneness for errors/accidents at work or while driving, tension headaches, and/or gastrointestinal symptoms associated with sleep loss or a concern about sleep). Patients were diagnosed with sleep apnea if PSGs showed an apnea hypopnea index (AHI; mean apnea/hypopneas or partial apneas per hour) >5, and they were diagnosed with PLMD if PSGs revealed myoclonus arousal >15/hour.

### Anxiety Measure

Anxiety was assessed through the State Trait Anxiety Inventory Form Y (STAI-Y). (Spielberger et al., 1983), a 20-item questionnaire measuring general levels of anxiety. Items consist of self-descriptive statements (e.g., “I feel \_\_\_\_; calm, secure, upset”) and participants are asked to rate their agreement on a 4-point Likert scale (1 = “not at all” to 4 “extremely”). Total scores range from 20–80, with higher scores indicating greater maladjustment. To match the two-week baseline period, STAI-Y instructions were modified to have participants respond based on the past two weeks. The STAI-Y demonstrates test-retest reliability exceeding .7 and reliably distinguishes patient and normal groups (Spielberger et al., 1983, Kabacoff et al., 1997).

### Cognitive Tasks

As part of the larger aims of this project, a computerized cognitive battery was administered immediately after the 14 days of actigraphy to examine cognitive functioning. From this larger battery, tasks were chosen to represent a range of cognitive performance in both lower order (i.e., attention/processing speed) and higher order (i.e., executive functioning) domains: Attention/Processing Speed [symbol digit modality task (SDMT), Simple Reaction Time (SRT)], Executive functioning (flanker task, letter series, N-back task).

**SDMT.**—The SDMT (Smith, 1982) measures sustained attention and processing speed (Spreen and Strauss, 1998). In this task, participants are presented with a legend at the top of the computer screen consisting of nine digit/symbol pairs. A series of symbols with blank spaces are shown below the legend. Participants are instructed to enter the corresponding number (provided in the legend) for each symbol, and to complete as many responses as

possible within 90 seconds. The number of correct responses entered in the time limit was computed (i.e., total items with correct numbers that correspond with the symbol), with higher scores representing better performance.

**SRT.**—In the SRT task, participants are presented with a warning stimulus (\*\*\*) followed by a signal stimulus (+) in the middle of the computer screen. Participants are instructed to press a designated key with their dominant hand as quickly as possible when the signal stimulus appears. Ten practice trials preceded the 50 test trials. Ten randomly arranged trials were then presented at each of five intervals separating the warning and signal stimuli (500, 625, 750, 875, and 1000ms). The main outcome measure was the average reaction time to respond to the 50 test trials, with higher scores indicating worse performance.

**N-Back.**—In this task, participants view a single letter in 48-point font in the center of the computer screen and decide whether that letter matched a target letter present N trials previously, with N varying from 0–2 trials. They are instructed to respond as fast and accurately as possible. In the 2-Back condition, participants judge whether the current letter matched the letter presented two letters previously, indicating their response with designated “yes”/“no” keyboard keys. Letters remained visible until a response was made, with a one second inter-stimulus interval. The N-Back task measures working memory, which is considered an executive function (Cohen et al., 1997). The outcome measure of interest was the total number of correct 2-back trials, with higher scores representing better performance.

### Letter Series.

The letter series test (Thurstone and Thurstone, 1962) measures inductive reasoning, an executive function. Participants are presented with a letter series on the computer screen and instructed to choose the letter (from five possible choices) that would continue the established pattern. Participants are to complete as many of the series as possible in four minutes. The proportion of correct trials (# correct/total trials attempted) was computed as the outcome measure of interest, where higher scores indicate better performance.

### Flanker Task.

The flanker task (Eriksen and Eriksen, 1974) involves responding to a target in the presence of non-target stimuli “flanked” around the target. Non-target stimuli are either congruent (i.e., correspond to the same directional response as the target), incongruent (i.e., correspond to the opposite response as the target), or neutral (i.e., do not correspond to any of the required responses regarding the target item). Our outcome of interest was the average reaction time for target response on incongruent trials. Performance on these trials measures the ability to inhibit irrelevant stimuli, an index of executive functioning (Eriksen and Eriksen, 1974). Higher scores indicate worse performance.

### Statistical Analyses

Multiple linear regressions were carried out using the PROCESS macro (Hayes, 2016) in SPSS (Version 24). Criterion variables included cognitive measures: SDMT, SRT, N-back task, letter series, flanker task. Predictor variables included: average TST (over 14 days of actigraphy), age, the TST by age interaction, and STAI-Y anxiety scores. Significant TST by

age interactions were clarified by calculating simple slopes of the association between TST and cognitive performance for sample-estimated age values characterized as follows: younger/middle-aged (1 SD below mean age, ~48 years of age), younger-older adults (~60 years of age) and older-older adults (1 SD above mean age, ~72 years of age).

## Results

### Participant Characteristics

Participant demographics, and values for sleep and cognitive measures are provided in Table 1. A total of 41 participants ( $M_{age}=60.3$ ,  $SD=12.3$ ) had full data available for actigraphy, anxiety, and cognition values and were included in analyses. Twelve participants (29%) met trial-defined criteria for insomnia, five (12%) participants met criteria for sleep apnea (AHI > 5), and no participants met criteria for PLMD.

### Multiple Regression

Results of models predicting performance for each cognitive test are presented in Table 2. Findings were similar for analyses including and excluding patients who met criteria for sleep apnea.

### Processing Speed/Attention Functioning.

For SDMT, TST and age independently predicted performance. Longer TST was associated with better performance, regardless of participant age. In general, older age was associated with worse performance. Anxiety scores were not associated with SDMT performance. For SRT, cognitive performance was not associated with any predictors.

### Executive Functioning.

TST independently predicted performance on letter series task and N-back tasks. Longer TST was associated with better letter series and N-back performance, regardless of participant age. Age and anxiety did not predict performance in either cognitive task.

Age independently predicted performance on the flanker task. Older age was associated with worse performance. TST and anxiety did not predict performance on the flanker task.

### Exploratory Analyses

To explore whether sleep fragmentation may have played a role in associations between TST and cognitive performance, we conducted additional separate multiple regressions, with WASO (Mean=50.2 minutes,  $SD=26.5$ , Range=5.9–144.8) and sleep efficiency percentage (SE; calculated as time spent asleep/time spent in bed x 100%; Mean=77.9,  $SD=9.5$ , Range=52.4–95.2), entered separately as predictors in the previously examined regression models. Results remained similar for associations between TST and performance across all tests. Further, WASO and SE did not predict performance on any test ( $ps >.05$ ).

## Discussion

The present study evaluated associations between objective TST, its interaction with age, as well as the independent association of anxiety, on cognitive performance in patients with ICDs and self-reported sleep disturbances. Regardless of age, TST predicted higher order cognition, and performance on a lower order cognitive task that measured processing speed and attention; whereas, anxiety was not associated with cognition.

Our prediction that TST would be associated with performance on tests measuring higher order cognitive functions was generally supported, but the type of cognitive task was important. For the reasoning and working memory tasks, associations between longer sleep duration and better performance agree with the direction of association reported in prior studies examining self-reported TST in adults without ICDs (Khasawneh et al., 2018, Fernandez-Mendoza et al., 2010). This suggests that patients with ICDs may receive cognitive benefits following increased sleep duration. The present results are generally consistent with prior research on older adults, which found similar associations with global cognitive performance for participants with long sleep duration (i.e., >8 hours; (Blackwell et al., 2011)). While we also predicted that TST associations would be strongest with increasing age, this was not supported. This may suggest that in patients with ICDs, the association between sleep duration and higher order cognition may be a generalized relationship across the lifespan.

Our prediction that TST would not be associated with lower level cognitive functioning in patients with ICDs was partially supported. The results showing no association between TST and SRT are not surprising given that sleep loss has been associated with the pre-frontal cortex (Horne, 1993), an area which is not likely to mediate this lower level cognitive task. Additionally, given that cognitive outcomes were assessed at single time-points, sleep duration may not have as much of an impact on cognitive performance as participants were likely able to recruit the necessary cognitive resources to perform this task. It is possible that examination of repeated daily cognitive assessments may reveal different associations with sleep duration than the single-shot approach reported here, important considerations for future research on patients with ICDs. However, we did observe that longer TST was associated with better SDMT performance. Given that SDMT measures attentional vigilance in addition to simple processing speed (Spren and Strauss, 1998, Shum et al., 1990), it presumably requires additional cognitive resources than those required to conduct SRT. Thus, like the other higher order cognitive tasks examined in the present study, SDMT may also be sensitive to disturbance following shorter sleep duration.

Interestingly, although all participants self-reported sleep disturbances, only a small proportion of patients met trial criteria for insomnia disorder or sleep apnea. Therefore, the present results demonstrate that even without criteria met for these sleep disorders, patients with ICDs still might experience impairments that limit their daytime functionality. Further, our finding that results remained similar whether patients with sleep apnea were or were not included in analyses agree with prior research showing associations between objectively measured (i.e., through PSG) sleep duration and cognitive performance in middle-aged adults with insomnia, independent of sleep disordered breathing (Fernandez-Mendoza et al.,

2010). Thus, in general, sleep duration may be an important area for clinical monitoring/target for intervention in patients with ICDs.

While not a primary focus of the present study, our exploratory analyses showing no associations between cognition and other objective sleep parameters that capture wake time during the night (i.e., WASO and SE), as well as no change in associations between TST and cognition, are interesting. While these exploratory analyses should be evaluated in larger samples, it is possible that longer sleep, regardless of how fragmented, is more important to improving higher order cognitive performance in patients with ICDs.

Contrary to our prediction, general anxiety scores were not associated with cognitive performance across any examined domain in patients with ICDs. These findings are surprising given the well-known negative association between anxiety and executive functions (Derakshan and Eysenck, 2009). We did not, however, examine associations between anxiety and all possible measures of executive function, but rather only on 3 tests measuring reasoning, working memory, and inhibition. Moreover, patients with ICDs may also have disease-specific anxiety, often termed *shock anxiety*, that could be more salient for this patient group (Ford et al., 2012). Therefore, it is possible that the pattern of results reported here may differ on other executive ability measures or different disease-specific anxieties. Further, given that recommended cutoff scores for clinically meaningful anxiety on the STAI-Y are higher for older adults, and the mean age of the present sample was on the older age range (i.e., ~60 years of age), anxiety scores in the present sample (mean of ~36) may not have been high enough to show the expected associations with higher order cognition. Finally, given that we measured anxiety and cognitive performance at single time-points, it is possible that there are acute fluctuations in the associations between anxiety and day-to-day cognitive performance in patients with ICDs, which should be examined in future work.

The present study has several limitations. First, while findings are likely representative of patients with ICDs and self-reported sleep disturbance, whether results may generalize to patients without self-reported sleep disturbances remains unclear. Second, the sample size was small; therefore, findings should be replicated in larger patient samples. Larger samples would also allow for sufficient power (adhering to the recommended sample size of 10 participants per predictor variable in the regression model (VanVoorhis and Morgan, 2007)) to explore other potential predictors (e.g., time since ICD implantation, medications, total number of shocks received) of cognitive performance. Third, as previously stated, given that we did not explore all possible test measures of each cognitive domain, it is possible that findings may not generalize to associations between sleep duration and performance on other cognitive tests. Fourth, given that the mean age of participants was approximately 60, results may not generalize to younger patients. Finally, as the present study did not compare findings to that of an age-matched control group of participants without ICDs, we cannot be certain that findings are specific to ICD patients.

To our knowledge, the present study is the first to examine associations between sleep and cognition in patients with ICDs. Results suggest that regardless of patient age, objectively assessed sleep duration may play an important role in the prediction of executive function



and a lower order task measuring attention and processing speed in patients with an ICD and self-reported sleep disturbance. Given that cognitive function is associated with quality of life in ICD patients (Moulaert et al., 2010), as well as functional independence in older adults (Mehta et al., 2002), who were widely represented in the present analysis, results may inform clinical applications related to sleep and aging, and potentially facilitate the management of cognitive aspects of daytime functioning.

## Acknowledgments

**Funding:** This project was supported by the National Heart, Lung, and Blood Institute (R21HL087831). Joseph M. Dzierzewski was supported by the National Institute on Aging of the National Institutes of Health (NIA; K23AG049955). The content is solely the responsibility of the authors and does not necessarily represent official views of the NIA.

## References

- Aasm. International classification of sleep disorders—third edition (ICSD-3). Darien, IL: American Academy of Sleep Medicine, 2014
- Association, A. P. Diagnostic and statistical manual of mental disorders (DSM-5®). American Psychiatric Pub, 2013 (
- Berg SK, Higgins M, Reilly CM, Langberg JJ and Dunbar SB Sleep quality and sleepiness in persons with implantable cardioverter defibrillators: outcome from a clinical randomized longitudinal trial. *Pacing Clin. Electrophysiol*, 2012, 35: 431–43. [PubMed: 22303998]
- Bilge AK, Ozben B, Demircan S, Cinar M, Yilmaz E and Adalet K Depression and anxiety status of patients with implantable cardioverter defibrillator and precipitating factors. *Pacing Clin. Electrophysiol*, 2006, 29: 619–26. [PubMed: 16784428]
- Blackwell T, Yaffe K, Ancoli-Israel S et al. Association of sleep characteristics and cognition in older community-dwelling men: the MrOS sleep study. *Sleep*, 2011, 34: 1347–56. [PubMed: 21966066]
- Cohen JD, Perlstein WM, Braver TS et al. Temporal dynamics of brain activation during a working memory task. *Nature*, 1997, 386: 604–8. [PubMed: 9121583]
- Derakshan N and Eysenck MW Anxiety, processing efficiency, and cognitive performance: New developments from attentional control theory. *Eur. Psychol*, 2009, 14: 168–76.
- Dornelas E and Sears SF Living with heart despite recurrent challenges: Psychological care for adults with advanced cardiac disease and arrhythmias. *Am. Psychol*, in press
- Edinger JD, Bonnet MH, Bootzin RR et al. Derivation of research diagnostic criteria for insomnia: report of an American Academy of Sleep Medicine Work Group. *Sleep*, 2004, 27: 1567–96. [PubMed: 15683149]
- Eriksen BA and Eriksen CW Effects of noise letters upon the identification of a target letter in a nonsearch task. *Percept. Psychophys*, 1974, 16: 143–49.
- Fernandez-Mendoza J, Calhoun S, Bixler EO et al. Insomnia with objective short sleep duration is associated with deficits in neuropsychological performance: a general population study. *Sleep*, 2010, 33: 459–65. [PubMed: 20394314]
- Ford J, Finch JF, Woodrow LK et al. The Florida Shock Anxiety Scale (FSAS) for patients with implantable cardioverter defibrillators: testing factor structure, reliability, and validity of a previously established measure. *Pacing Clin. Electrophysiol*, 2012, 35: 1146–53. [PubMed: 22758923]
- Habibovi M, Mudde L, Pedersen S, Schoormans D, Widdershoven J and Denollet J Sleep disturbance in patients with an implantable cardioverter defibrillator: Prevalence, predictors and impact on health status. *Eur. J. Cardiovasc. Nurs*, 2018: 1474515117748931.
- Hałas K, Krzy anowski K, Krzy anowska E et al. Cognitive impairment after appropriate implantable cardioverter-defibrillator therapy for ventricular fibrillation. *Kardiologia Polska (Polish Heart Journal)*, 2014, 72: 134–39. [PubMed: 23990235]

- Hayes AF The PROCESS macro for SPSS and SAS. URL: <http://www.processmacro.org/index.html> (visited on 12/07/2017), 2016
- Heron MP Deaths: Leading causes for 2015. 2017
- Horne JA Human sleep, sleep loss and behaviour: implications for the prefrontal cortex and psychiatric disorder. *The British Journal of Psychiatry*, 1993, 162: 413–19. [PubMed: 8453439]
- Kabacoff RI, Segal DL, Hersen M and Van Hasselt VB Psychometric properties and diagnostic utility of the Beck Anxiety Inventory and the State-Trait Anxiety Inventory with older adult psychiatric outpatients. *J. Anxiety Disord*, 1997, 11: 33–47. [PubMed: 9131880]
- Kamphuis HC, Verhoeven NW, De Leeuw R, Derksen R, Hauer RN and Winnubst JA ICD: a qualitative study of patient experience the first year after implantation. *J. Clin. Nurs*, 2004, 13: 1008–16. [PubMed: 15533107]
- Khassawneh BY, Bathgate CJ, Tsai SC and Edinger JD Neurocognitive performance in insomnia disorder: The impact of hyperarousal and short sleep duration. *J. Sleep Res*, 2018: e12747. [PubMed: 30069961]
- Kusumoto FM, Calkins H, Boehmer J et al. HRS/ACC/AHA expert consensus statement on the use of implantable cardioverter-defibrillator therapy in patients who are not included or not well represented in clinical trials. *J. Am. Coll. Cardiol*, 2014, 64: 1143–77. [PubMed: 24820349]
- Lichstein KL, Durrence HH, Taylor DJ, Bush AJ and Riedel BW Quantitative criteria for insomnia. *Behav. Res. Ther*, 2003, 41: 427–45. [PubMed: 12643966]
- Lichstein KL, Riedel BW and Means MK Psychological treatment of late-life insomnia. 1999
- Lichstein KL, Stone KC, Donaldson J et al. Actigraphy validation with insomnia. *Sleep*, 2006, 29: 232–9. [PubMed: 16494091]
- Lo JC, Loh KK, Zheng H, Sim SK and Chee MW Sleep duration and age-related changes in brain structure and cognitive performance. *Sleep*, 2014, 37: 821–21. [PubMed: 24899768]
- Mccrae CS, Roth AJ, Ford J et al. Sleep, psychosocial functioning, and device-specific adjustment in patients with implantable cardioverter defibrillators (ICDs). *Behav. Sleep Med*, 2016, 14: 49–66. [PubMed: 25174823]
- Mccrae CS, Rowe MA, Tierney CG, Dautovich ND, Definis AL and Mcnamara JP Sleep complaints, subjective and objective sleep patterns, health, psychological adjustment, and daytime functioning in community-dwelling older adults. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 2005, 60: P182–P89.
- Mehta KM, Yaffe K and Covinsky KE Cognitive impairment, depressive symptoms, and functional decline in older people. *J. Am. Geriatr. Soc*, 2002, 50: 1045–50. [PubMed: 12110064]
- Moulaert VR, Wachelder EM, Verbunt JA, Wade DT and Van Heugten CM Determinants of quality of life in survivors of cardiac arrest. *J. Rehabil. Med*, 2010, 42: 553–58. [PubMed: 20549160]
- Oakley N Validation with polysomnography of the Sleepwatch sleep/wake scoring algorithm used by the Actiwatch activity monitoring system. Bend: Mini Mitter, Cambridge Neurotechnology, 1997
- Pedersen SS, Van Den Broek KC, Theuns DA et al. Risk of chronic anxiety in implantable defibrillator patients: a multi-center study. *Int. J. Cardiol*, 2011, 147: 420–23. [PubMed: 19896732]
- Salthouse TA When does age-related cognitive decline begin? *Neurobiol. Aging*, 2009, 30: 507–14. [PubMed: 19231028]
- Sears SF, Hauf JD, Kirian K, Hazelton G and Conti JB Posttraumatic stress and the implantable cardioverter-defibrillator patient: what the electrophysiologist needs to know. *Circulation: Arrhythmia and Electrophysiology*, 2011, 4: 242–50. [PubMed: 21505176]
- Shum DH, Mcfarland KA and Bain JD Construct validity of eight tests of attention: Comparison of normal and closed head injured samples. *The Clinical Neuropsychologist*, 1990, 4: 151–62.
- Smith A Symbol digit modalities test (SDMT) manual (revised) Western psychological services. Los Angeles, 1982
- Sola CL and Bostwick JM Implantable cardioverter-defibrillators, induced anxiety, and quality of life (In). *Mayo Clin. Proc Elsevier*, 2005: 232–37.
- Spielberger CD, Gorsuch RL, Lushene R, Vagg PR and Jacobs GA State-Trait Anxiety Inventory, Form Y. Consulting Psychologists Press, Palo Alto, CA, 1983 (

Spree O and Strauss E A compendium of neuropsychological tests: Administration, norms, and commentary. Oxford University Press, 1998 (

Thurstone TG and Thurstone LL Primary mental abilities tests. Science Research Associates, 1962 (

Vanvoorhis CW and Morgan BL Understanding power and rules of thumb for determining sample sizes. Tutorials in Quantitative Methods for Psychology, 2007, 3: 43–50.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 1**

Demographics and sleep outcomes of study sample of ICD patients with self-reported sleep disturbance ( $N=41$ )

Variable	Mean (SD)	Range
Age	60.34 (12.32)	20.00–84.00
Sex (M:F)	23:18	--
Time since ICD implantation (years)	3.61 (3.14)	(0.08–13.00)
Medical Conditions		
Coronary Artery Disease (n,%)	7 (17%)	--
Cardiomyopathy (n, %)	3 (7%)	--
Congestive Heart Failure (n,%)	14 (34%)	--
History of Myocardial Infarction (n,%)	6 (15%)	--
Atrial Fibrillation (n,%)	19 (46%)	--
Psychiatric Diagnosis (n,%)	12 (29%)	--
Use of prescription sleep meds (n, %)	26 (63%)	--
MMSE	27.11 (1.82)	26.00–30.00
Anxiety score (STAI-Y)	36.15 (13.35)	20.00–72.00
<i>Sleep Measure</i>		
TST-actigraphy (minutes)	409.77 (74.77)	262.20–566.43
<i>Cognitive Measures</i>		
Flanker Task (RT in seconds)	710.78 (207.00)	227.40–1184.63
Letter Series (proportion correct)	.59 (.29)	.06–1.00
N-Back (2-back; number correct)	83.30 (19.11)	16.00–100.00
SDMT (number correct)	28.32 (11.36)	1.00–56.00
SRT (RT in seconds)	344.87 (121.21)	182.43–673.74

*Note.* Dx = diagnosis; MMSE = Mini-Mental State Examination; STAI-Y = State-Trait Anxiety Subscale Form Y; TST = total sleep time; SDMT = Symbol Digit Modality Test; SRT = Simple Reaction Time

**Table 2**

Actigraphic sleep determinants of cognitive performance in patients with an Implantable Cardiac Defibrillators (ICD) ( $n=41$ )

Cognitive Domain/Measure	<i>B</i>	<i>SE</i>	<i>t</i>	<i>p</i>
<i>Attention/Processing Speed</i>				
<b>SDMT<sup>a</sup></b>				
TST	.05	.02	2.37	.02
Age	-.42	.14	-2.91	.01
TST x age	-.00	.00	-.85	.40
Anxiety score	-.11	.13	-.82	.42
<b>SRT<sup>a</sup></b>				
TST	-.29	.24	-1.17	.25
Age	2.85	1.57	1.81	.08
TST x age	.01	.03	.28	.78
Anxiety score	-2.79	1.46	-1.91	.07
<i>Executive Functioning</i>				
<b>N-Back (2-back)<sup>a</sup></b>				
TST	.10	.04	2.53	.02
Age	-.08	.26	-.33	.74
TST x age	-.01	.00	-1.44	.16
Anxiety score	.03	.24	.14	.89
<b>Letter Series<sup>a</sup></b>				
TST	.00	.00	2.19	.04
Age	-.00	.00	-1.10	.28
TST x age	-.00	.00	-.63	.53
Anxiety score	.00	.00	.34	.74
<b>Flanker Task<sup>a</sup></b>				
TST	-.38	.40	-.96	.34
Age	8.81	2.56	3.43	<.01
TST x age	-.02	.05	-.35	.73
Anxiety score	-.87	2.39	-.36	.72

Note. *SE* = Standard error; Flanker Task = Mean reaction time on incongruent trials;

TST = total sleep time; Anxiety score = Score on the STAI-Y; SDMT = Symbol Digit Modality Test;

SRT = Simple Reaction time.

<sup>a</sup>Regression analyses were also conducted with participants who met sleep apnea criteria (AHI > 5,  $n = 5$ ) removed. Results remained similar across all cognitive tasks, therefore these participants were retained in the study sample.